

The client sample was analyzed for plant-based cannabinoids by Liquid Chromatography (LC). The collected data was compared to data collected for certified reference standards at known concentrations.

70459-CN

ID	Weight %	Concentration (mg/mL)			
D9-THC	ND	ND			
THCV	ND	ND			
CBD	9.82	95.46			
CBDV	ND	ND			
CBG	ND	ND			
CBC	ND	ND			
CBN	ND	ND			
THCA	ND	ND			
CBDA	ND	ND			
CBGA	ND	ND			
D8-THC	ND	ND			
exo-THC	ND	ND			
Total	9.82	95.46	0%	Cannabinoids (wt%)	9.8%
Max THC	ND	ND			
Max CBD	9.82	95.46			

Limit of Quantitation (LOQ) = 0.01 wt%

Max THC (and Max CBD) are calculated values for total cannabinoids after heating, assuming complete decarboxylation of the acid to the neutral form. It is calculated based on the weight loss of the acid group during decarboxylation: Max THC = $(0.877 \times THCA) + THC$. This calculation does not include other cannabinoid isomers (eg. D8-THC and exo-THC). ND = None detected above the limits of detection (LOD), which is half of LOQ.

TP: Terpenes Profile [WI-10-27]Analyst: JRTest Date: 11/12/2019

The client sample was analyzed by Head-Space Gas Chromatography (HS-GC). The collected data was compared to data collected for certified reference standards at known concentrations. All values are semiquantitative estimates based on recorded peak areas relative to terpene calibration data.

70459-TP

	Compound	ppm	Terpene	Profile		Compound	ppm	Terpene	e Profile	
	beta-myrcene					camphene				
	isopulegol					L-fenchone				
	menthol					beta-pinene				
	cis-nerolidol					eucalyptol				
	trans-nerolidol					alpha-terpinene				
	gamma-terpinene					delta-3-carene				
	alpha-bisabolol					alpha-pinene				
	linalool					D-limonene				
b	eta-caryophyllene					geraniol				
ca	ryophyllene oxide					cis-beta-ocimene				
	guaiol					alpha-ocimene				
	sabinene					alpha-phellandrene				
	alpha-humulene					terpinolene				
	p-cymene									
Tot	pp al Terpene: <0	om 0.0 .1 wt%		00	10.00		0.0	0 5.	00	10.00

END OF REPORT